

U. S. Geological Survey
Geologic Branch Library

USGS LIBRARY-DENVER



3 1819 00045426 0

THE

AMERICAN JOURNAL

OF

SCIENCE AND ARTS.

EDITORS,

JAMES D. DANA, B. SILLIMAN, AND E. S. DANA.

ASSOCIATE EDITORS,

PROFESSORS ASA GRAY, WOLCOTT GIBBS AND
J. P. COOKE, JR. OF CAMBRIDGE,

PROFESSORS H. A. NEWTON, W. JOHNSON, G. J. BRUSH
AND A. E. VERRILL OF NEW HAVEN,

PROFESSOR GEORGE F. HARKER, OF PHILADELPHIA.

THIRD SERIES.

VOL. XV.—[WHOLE NUMBER, CXV.]

Nos. 85—90.

JANUARY TO JUNE, 1878.

WITH TWO PLATES.

4799

NEW HAVEN: EDITORS.

1878.

NUMBER LXXXIX.

	Page
ART. XLVII.—Research on the Absolute Unit of Electrical Resistance; by HENRY A. ROWLAND,	325
XLVIII.—Meteoric Iron from Virginia; by J. W. MALLETT, ..	337
XLIX.—Relative Positions of the Forest Bed and associated Drift Formations in Northeastern Iowa; by W. J. MCGEE, ..	339
L.—Geographical and Geological Survey of the Rocky Mountain Region, under the direction of J. W. POWELL, ..	342
LI.—Just Intonation in Music; by H. WARD POOLE,	359
LII.—Brachiopoda in the Swedish Primordial; by S. W. FORD, ..	364
LIII.—On the "Geodes" of the Keokuk Formation, and the Genus Biopalla; by SAMUEL J. WALLACE, ..	366
LIV.—The Coralline, or Niagara Limestone of the Appalachian System; by S. T. BARRETT,	370
LV.—Isopoda from New England; by OSCAR HARGER,	373
LVI.—Ammonio-argentic Iodide; by M. CAREY LEA,	379
LVII.—Fossil Passerine Bird from Colorado; by J. A. ALLEN, ..	381
LVIII.—Terrestrial Electrical Currents; by W. L. BROWN, ..	385
LIX.—Notice of New Fossil Reptiles; by O. C. MARSH,	409

SCIENTIFIC INTELLIGENCE.

Chemistry and Physics.—Specific Heat of Beryllium, NILSON and PETTERSSON: Synthesis of Olefines, ELTEKOFF, 386.—Transformation of Olefine bromides into bromides of radicals of the Fatty Acids, DEMOLE: Reaction of Boric acid and Borax, etc., KLEIN, 387.—Distillation of Resins and resin acids with zinc dust, CIAMICIAN: Triacid Phenols of Wood Tar, etc., HOFMANN, 388.—Constituents of Corallin, ZULKOWSKY: Curarine, SACHS: Protection of Marine Boilers, 389.—Boric Acid, M. A. DITTE, 390.—Boiling Point of Iodide of Antimony, 391.—Influence of the density of a body upon its Light-absorbing power, GLAN: Reflection of Polarized Light from the Equatorial surface of a Magnet, KERR: Method of measuring the Velocity of Light, A. A. MICHELSON, 394.

Geology and Mineralogy.—Limestones of the Falls of the Ohio, JAMES HALL, 395.—Fossil Plants of the Auriferous Gravel Deposits of the Sierra Nevada, L. LESQUEREUX: Fossil wood from the Keokuk formation, S. J. WALLACE: Atlas accompanying the Report of the Geological Exploration of the Fortieth Parallel. C. KING, 396.—Geological and Geographical Atlas of Colorado, F. V. HAYDEN: Genera Zethus, Cybele, Encrinurus and Cryptonotus, A. W. VOGES, 397.—Carboniferous and Upper Silurian fossils of Illinois and Indiana, C. A. WHITE: Culm-Flora der Ostrauer und Waldenburger Schichten, D. STUR: Remains of the Musk Ox: Three new Phosphates from Fairfield County, Conn., G. J. BRUSH and E. S. DANA, 398.—General Principles of Mineralogy, J. H. COLLINS, 399.

Botany and Zoology.—Synoptical Flora of North America, GRAY: Bibliographical Index to North American Botany, S. WATSON, 400.—Some points in the Morphology of Primulaceæ, M. T. MASTERS: Origin of Floral Æstivations, etc., G. HENSLow, 401.—Floral Structure and Affinities of Sapotaceæ, M. M. HARTOG: North American Plants: Spore-Formation of the Mesocarpeæ, V. B. WITTROCK, 402.—Non-Sexual Outgrowths on Fern Prothalli, 403.—Catalogue of the Flowering Plants within thirty miles of Yale College, 404.

Astronomy.—Der Sternhaufen χ Persei, etc., H. C. VOGEL, 404.—American Journal of Mathematics: Publications of the Cincinnati Observatory, 406.

Miscellaneous Scientific Intelligence.—Driftless Region of Wisconsin, 406.—Treatise on Chemistry, H. E. ROSCOE and C. SCHORLEMMER: Report of the Superintendent of the Coast Survey for 1874: Minnesota Academy of Natural Sciences: Matter and Motion, J. C. MAXWELL: Contributions to North American Ethnology, S. POWERS, 407.—Obituary.—CHARLES PICKERING, 408.

ART. LVII.—Description of a Fossil Passerine Bird from the Insectbearing Shales of Colorado; by J. A. ALLEN.*

THE species of fossil bird described in this paper is based on some beautifully preserved remains from the insect-bearing shales of Florissant, Colorado. They consist of the greater part of a skeleton, embracing all of the bones of the anterior and posterior extremities, excepting the femora. Unfortunately, the bill and the anterior portion of the head are wanting, but the outlines of the remainder of the head and of the neck are distinctly traceable. The bones are all *in situ*, and indicate beyond question a high ornithic type, probably referable to the Oscine division of the *Passeres*. The specimen bears also remarkably distinct impressions of the wings and tail, indicating not only the general form of these parts, but even the shafts and barbs of the feathers.

In size and in general proportions, the present species differs little from the Scarlet Tanager (*Pyrrhula rubra*) or the Cedar-bird (*Ampelis cedrorum*).† The bones of the wings, as well as the wings themselves, indicate a similar alar development, but the tarsi and feet are rather smaller and weaker; and hence in this point the agreement is better with the short-legged Pewees (genus *Contopus*). These features indicate arboreal habits and well-developed powers of flight. The absence of the bill renders it impossible to assign the species to any particular family, but the fossil on the whole gives the impression of Fringilline affinities.

Palæospiza bella, gen. et sp. nov.

Wings rather long, pointed. Tail (apparently)† about two-thirds the length of the wing, rounded or graduated, the outer feathers (as preserved) being much shorter than the inner. One side shows distinctly six rectrices. Tarsus short, its length a little less than that of the middle toe. Lateral toes subequal, scarcely shorter than the middle one. Hind toe about two-thirds as long as the middle toe. Feet and toes strictly those of a perching bird, and the proportionate length of the bones of the fore and hind limbs is the same as in ordinary arboreal *Passeres*, especially as represented by the *Tanagridæ*.

* From the Bulletin of the Geological and Geographical Survey of the Territories, vol. iv, No. 2, page 443, April, 1878.

† The character of the tail is given with reservation, since it is not quite certain that the whole of the tail, or that the exact form of the terminal portion, is shown, especially as the preserved impression is somewhat unsymmetrical.



One of the specimens affords the following measurements :

	Inches.		Inches.
Humerus, length.....	0.80	Middle toe and claw.....	0.65
Forearm, length.....	0.95	Claw alone.....	0.20
Manus, length.....	1.02	Hind toe and claw.....	0.37
Coracoid, length.....	0.72	Claw alone.....	0.15
Clavicle, length.....	0.63	Wing.....	3.60
Tibia, length.....	1.00	Tail (approximate).....	2.70
Tarsus, length.....	0.60	Total length (approximate).....	6.85

The bones still rest in the original matrix, and, being somewhat crushed and flattened, do not admit of detailed description and comparison with other types. The furculum is well preserved, and the limb-bones are all in place in their natural relation. The sternum is unrecognizable. The position of the cervical series of vertebræ and the general outline of the skull can be traced; but no structural characters of the head can be distinguished, except the proximal portion of the mandible. The long bones all present a well-marked longitudinal groove, due evidently to compression and fracture. This groove is distinctly traceable, even in such slender bones as tibiæ, tarsi, and clavicles. In point of size, while the furculum and the bones of the wing have all about the same length as the corresponding parts in *Ampelis cedrorum*, they apparently are considerably stouter. Their greater breadth may, however, be due simply to flattening from pressure. The tibiæ and tarsi are a little shorter than in the species last named, but the difference is only slight.


The most remarkable feature of the specimen is the definiteness of the feather impressions. Both the shafts and the barbs are shown with great distinctness in the rectrices, and the tips of the primaries of one wing are also sharply defined, overlying the edge of the partly expanded tail. The tip of the opposite wing can also be seen beneath the tail. The feet are so beautifully preserved that even the claws are perfectly distinct. (Fig. 1.)

Another specimen from the same locality, and probably representing the same species, consists of the tip of the tail and about the apical third of a half-expanded wing. (Fig. 2.) In this example the tail is also pointed and graduated. About seven of the outer primaries of the wing are shown with great distinctness, and two others can be easily made out. The third primary is the longest; the second is slightly shorter; the first and fourth are about equal. There are also in the collection three detached contour feathers of small size, but whether pertaining to the same species as the other specimens cannot, of course, be determined.

The larger specimen, first described, is divided into an upper and a lower half, the greater part, however, adhering to

the lower slab. The bones adhere about equally to the two faces. The drawing is made from the lower slab, with some of the details filled in from the upper one. The feather impressions are about equally distinct on both, and where in either case the bones are absent, exact molds of them remain, so that the structure can be seen and measurements taken almost equally well from either slab, except that nothing anterior to the breast is shown on the upper slab.

2.



The species here described is of special interest as being the first fossil Passerine bird discovered in North America, although birds of this group have been known for many years from the Tertiary deposits of Europe. The highest extinct ornithic type hitherto known from America is a Picarian bird (*Uintornis lucaris*) related to the Woodpeckers, described by Professor O. C. Marsh in 1872, from the Lower Tertiary of Wyoming Territory. Probably the insect-bearing shales of Colorado will afford, on further exploration; other types of the higher groups of birds.

For the opportunity of describing these interesting specimens I am indebted to Mr. S. H. Scudder, who obtained them during his last season's (1877) explorations of the Florissant insect-beds. The specimens are now the property of the Boston Society of Natural History. My thanks are due to Mr. J. H. Blake for the great care with which he has executed the drawings.

In conclusion, I may add that in 1871 I obtained a few distinct impressions of feathers from beds of the same age and from near the same locality. The first fossil feather, to my knowledge, discovered in North America, was obtained by Dr. F. V. Hayden in 1869, from the fresh-water Tertiary deposits of Green River, Wyoming Territory. This was described by Professor O. C. Marsh in 1870,* who refers to it as "the distal portion of a large feather, with the shaft and vane in excellent preservation."

* This Journ., II, vol. xi, p. 272, 1870.